

farsida 8 juni 2022

by Farsida 8 Juni 2022

Submission date: 08-Jun-2022 12:42PM (UTC+0700)

Submission ID: 1852759629

File name: Farsida_Final_8.06.2022.docx (111.18K)

Word count: 5032

Character count: 26836

FACTORS ASSOCIATED WITH BCG SCAR OF PEDIATRIC TUBERCULOSIS PATIENTS AT PISANGAN AND EAST CIPUTAT COMMUNITY HEALTH CENTERS

Farsida^{1*}, Ayu Farah Syifa¹, Azka Zifa Tanama¹

¹ Faculty of Medicine and Health, Universitas Muhammadiyah Jakarta

*Correspondence email: farsida@umj.ac.id

ABSTRACT

Tuberculosis (TB) is one of the most common causes of death in children. Prevention of TB through Bacille Calmette–Guérin (BCG) vaccination is an action to produce immunity against TB disease. After vaccination, a scar forms from the boil, which indicates a successful vaccination. This search aims to determine the factors that can be associated with BCG Scar in pediatric TB patients. A cross-sectional design was conducted at the Pisangan and East Ciputat Community Health Centers in January–September 2020. Data collection using medical records and telephone interviews on 35 samples was taken using the probability sampling technique, simple random sampling. With inclusion criteria for pediatric TB patients (0-18 years) and exclusion criteria that do not include telephone numbers and addresses. Bivariate analysis was used with the chi-square test. The results showed a significant relationship between contact history and BCG scars in pediatric TB patients. Age, gender, nutritional status, exclusive breastfeeding, immunization, mother's education, parent's occupation, family income, and smoking history were not significantly related to BCG scar. Contact history related to BCG scar. Parents should be giving more attention to and protecting their children from the people around them to cut off the chance of contact with TB sufferers even though the child has been vaccinated.

Keywords: Child TB; Contact History; Scar BCG

INTRODUCTION

Tuberculosis (TB) is one of the infectious diseases that has become the leading to ill health. TB is also becoming one of the top 10 diseases that cause death globally. Moreover, TB got ranked above HIV/AIDS as the cause of death from a single infectious agent. The bacillus *Mycobacterium tuberculosis* is the cause that made TB happen, and this bacillus spreads every time a person with TB expels bacteria into the air, especially when they cough. Globally in 2019, an estimated 10.0 million people in the range of 8.9-11.0 had TB. This number has declined quite slowly in recent years. In 2019 accounted 12% of children (aged <15 years) developed TB. Indonesia occupies the second position with the highest TB cases globally, with 543,874 cases in 2019.¹ In the same year

in Banten province, there were 3,123 TB cases in children 4 years old.²

The Bacille Calmette–Guérin (BCG) vaccine is one of the common interventions to control and prevent TB infection in children. This vaccination can give protection in children from severe forms of TB. A scar will form after the BCG vaccination, which indicates a successful BCG vaccination. If a boil doesn't form, it doesn't mean that the BCG vaccine didn't work or protection didn't form, so there's no need to repeat it even if a boil or scar doesn't form.³

Children vaccinated with BCG and had scars overall had better survival than children vaccinated with BCG and had no scars.⁴ One important marker for showing a properly functioning immune system is scar BCG. Inexpensive and direct initiatives such as early BCG vaccination, BCG scar tissue

monitoring, and revaccination of scar-negative children can directly impact child survival worldwide.⁵

Children who have scars are said to have fewer skin infections and se¹²s, and this is because BCG is a type of non-specific immune stimulator. It is confirmed by recent research that BCG produces a Th1 immune response at birth. ¹²⁵s also associated with mortality, which is 45-55% lower during the first year and a half of life than those without scars.¹

There is a relationship between the incidence of TB infection and the BCG vaccine scar formation. In Rosandali's study, people who did not have BCG scars were more common⁵ in TB patients.⁷ BCG scar readings have been used as an indicator of vaccine status and are becoming standard practice for assessing the protective effect of BCG in retrospective studies. A BCG scar has been an indication of protection against TB. However, more than fifty percent of the research population with BCG ³ars were infected with TB.⁵ Based on this, researchers are interested in knowing the factors associated with BCG Scar in pediatric TB patients.

MAT¹⁰RIAL AND METHODS

This stud²⁴ used a descriptive type of research using a cross-sectional design. This research was conducted at the Pisangan and East Ciputat Health Centers from January to September²⁸ r 2020. The sample was selected using a probability sampling technique of simple r¹¹om sampling with a total of 35 samples. The number of samples is calculated by the Slovin formula:⁸

$$n = \frac{N}{1 + Nd^2}$$

Inclusion criteria in this study were ¹⁹diatric TB patients (aged 0-18 years) and willing to participate in the study. The exclusion criteria are respondents who do not include telephone numbers and addresses. The research instrument used a questionnaire to obtain data on age, gender, TB contact history, nutritional status, immunization status, history of exclusive breastfeeding,

BCG vaccination scars, economic status, mother's education, parent's occupation, smoking history in parents taken via phone. Assessment of nutritional status based on weight indicators for age (weight/age). The secondary data used are medical records of pediatric patients with TB who have been ²⁶ccinated against BCG. The data were with univariate and bivariate analysis with a chi-square test, which was processed using SPSS 21.0 program.

RESULT

This study was conducted on 35 respondents of children aged 0-18 years with a diagnosis of TB who were treated with OAT at the Pisangan and East Ciputat³ Health Centers. The results showed that the characteristics of respondents based on age, children aged > 6-18 years were more (60.0%) compared to those aged six years, 62.9% were male and 37.1% female. The data also showed as many as 48.6% of respondents had normal nutritional status, 28.6% were undernourished, 5.7% were malnourished and overweight, and 11.4% were obese. Respondents with a history of exclusive breastfeeding for six months were 82.9%, most of the respondents had received complete immunization as many as 82.9%, respondents had a history of household contact as much as 34.3%, 22.9% not at home and unknown contacts 42.9%. Respondents with positive BCG scars were more (80.0%) than negative BCG scars (Table 1).

Characteristics of parents obtained that as many as 62.9% have higher education, mothers who do not work as much as 77.1% more than mothers who work, fathers who work as laborers 42.9% more than work as employees 20.0% and self-employed 37.1%, family income 62.9% have income \geq minimum wage (UMP). Also, parents have a history of smoking is high²⁰ than those not smoking as much as 62.9% (Table 1).

Table 1. Child and Parental Characteristics (n=35)

Variable	N	%
Child Characteristics		
Age		
≤ 6 years	14	40.0
> 6-18 years	21	60.0
Sex		
Boys	22	62.9
Girls	13	37.1
Nutrition Status		
Several underweight	2	5.7
Underweight	10	28.6
Normal	17	48.6
Overweight	2	5.7
Obesity	4	11.4
Exclusive Breastfeeding History		
< 6 months	6	17.1
≥ 6 months	29	82.9
Immunization Status		
Incomplete	6	17.1
Complete	29	82.9
Contact History		
Household	12	34.3
Not at home	8	22.9
Unknown	15	42.9
Scar BCG		
Negative	7	20.0
Positive	28	80.0
Parental Characteristics		
Mother's Education		
Low	13	37.1
High	22	62.9
Mother's Occupation		
Employee	8	22.9
Housewife	27	77.1
Father's Occupation		
Laborer	15	42.9
Employee	7	20.0
Entrepreneur	13	37.1
Family Income		
< Minimum wage (UMP)	13	37.1
≥ Minimum wage (UMP)	22	62.9
Smoking history		
Yes	22	62.9
No	13	37.1

31

Based on the results of bivariate analysis (Table 2), children aged six years with negative BCG scars (42.9%) and positive BCG scars (39.3%) were fewer than children aged > 6-18 years. The analysis results

showed no significant relationship between the child's age and the BCG scar (p-value 0.594). Girls had more negative BCG scars (57.1%) than boys (42.9%), but boys had more positive BCG scars (67.9%) than girls (32, 1%). The results of the chi-square analysis showed that there was no significant relationship between sex and BCG scars (p-value 0.214). Children with underweight have more negative BCG scars (57.1%) than children with normal nutrition (42.9%). Normal status has more positive BCG scars (71.4%) than children with underweight (28.6%). The results of the chi-square analysis showed that there was no significant relationship between nutritional status and BCG scars (p-value 0.163).

Children with a history of exclusive breastfeeding had a higher negative BCG scar (85.7%) than non-exclusive (14.3%), as well as a positive BCG scar in children with a history of exclusive breastfeeding, which was higher (82.1%) than non-exclusive (17.9%). The results of the chi-square analysis showed that there was no significant relationship between a history of exclusive breastfeeding and BCG scars (p-value 0.656). Children with complete immunization had negative BCG scars as many as seven people (100.0%), while children with incomplete immunizations did not have negative BCG scars. Children who were fully immunized had more positive BCG scars (78.6%) than children who were not fully immunized (21.4%). The results of the chi-square analysis showed that there was no significant relationship between immunization status and BCG scar (p-value 0.232). Respondents with a history of household contact as 42.9% and 57.1% did not have a negative BCG scar. Respondents with positive BCG scars who account for household contact are 32.1%, 14.3% are not at home, and 53.6% are unknown contacts. The results of the chi-square analysis showed that there was a significant relationship between contact history and BCG scars (p-value 0.015). Based on the characteristics of parents, mothers with higher education have more children with

negative BCG scars (57.1%) than mothers with low education (42.9%).

Table 2. Factors associated with Scar BCG

Variable	Scar BCG (-)		Scar BCG (+)		p-value	OR	CI 95%
	N	%	N	%			
Age					0.594	1.159	0.216-6.207
≤ 6 years	3	42.9	11	39.3			
> 6-18 years	4	57.1	17	60.7			
Sex					0.214	0.400	0.065-1.933
Boys	3	42.9	19	67.9			
Girls	4	57.1	9	32.1			
Nutrition Status					0.163	3.333	0.605-18.371
Underweight	4	57.1	8	28.6			
Normal	3	42.9	20	71.4			
Exclusive Breastfeeding History					0.656	0.767	0.075-7.860
< 6 months	1	14.3	5	17.9			
≥ 6 months	6	85.7	23	82.1			
Immunization Status					0.232		
Incomplete	0	0.0	6	21.4			
Complete	7	100.0	22	78.6			
Contact History					0.015*		
Household	3	42.9	9	32.1			
Not at home	4	57.1	4	14.3			
Unknown	0	0.0	15	53.6			
Mother's Education					0.525	1.35	0.250-7.278
Low	3	42.9	10	35.7			
High	4	57.1	18	64.3			
Mother's Occupation					0.516	1.467	0.226-9.534
Employee	2	28.6	6	21.4			
Housewife	5	71.4	22	78.6			
Father's Occupation					0.219		
Laborer	5	71.4	10	35.7			
Employee	1	14.3	6	21.4			
Entrepreneur	1	14.3	12	42.9			
Family Income					0.475	0.618	0.102-3.765
< UMP	2	28.6	11	39.3			
≥ UMP	5	71.4	17	60.7			
Smoking history					0.475	1.618	0.266-9.852
Yes	5	71.4	17	60.7			
No	2	28.6	11	39.3			

More children with mothers with higher education (64.3%) than children with low maternal education¹ (35.7%) with positive BCG scars. The results of the chi-square analysis showed that there was no significant relationship between maternal education and BCG scar (p-value 0.525). Mothers who do not work have more children with negative BCG scars (71.4%) than work³³ mothers (28.6%). Likewise, the positive BCG scar of the children with mothers who do not come to work is higher (78.6%)³ than children with mothers who work (35.7%). The results of the chi-square analysis showed that there was no

significant relationship between the mother's occupation and BCG scar (p-value 0.516). Fathers who work as laborers have more children with negative BCG scars (71.4%) than fathers who work as employees and entrepreneurs. Meanwhile, fathers who work as entrepreneurs have children with positive scars (42.9%) more than those who work as laborers and employees. The results of the chi-square analysis showed that there was no significant relationship between the father's occupation and BCG scar (p-value 0.219). Families with income ≥UMP have more children with negative BCG scars (71.4%)

than income <UMP (28.6%). Likewise, positive BCG scars for families with income \geq UMP more (60.7%) than income <UMP (39.3%). Parents who have a history of smoking have children with more negative BCG scars (71.4%) than parents who do not have a history of smoking (28.6%). Likewise, the positive BCG scars of parents who had a history of smoking were higher (60.7%) than parents who had no history of smoking (39.3%) (Table 2).

DISCUSSION

Child Characteristic

In this study, the age of respondents with the highest incidence of TB were respondents aged six years and over. In contrast to the research by Brajadenta et al. (2018), from 160 children diagnosed with TB, 83 children were aged 0-5 years, while other children were aged 5-14 years. In Indonesia, TB is one of the most common causes of mortality and morbidity in children. Children are very susceptible to TB infection. Children aged five years have a greater risk of developing TB infection because of their cellular immunity in their age is not fully developed (immature).

A cellular adaptive immune response provided by the ability of *Mycobacterium tuberculosis* to survive intracellularly. It is played by specific T cells in order to defend against TB. B cells, T Follicular B helpers, T regulators, restriction CD1, CD6+, CD4+, Th1, Th2, and Th17 are giving response to Bacillus of Calmette and Guerin (BCG) immunization as the adaptive immune response. Current childhood TB infection represents a future source of TB disease.

There are more TB patients in boys than girls. This study is in line with previous research, which stated that the gender of children with TB was more common in boys than girls. This is different from the research conducted by Permatasari and Trijati (2014), which stated that the sex research results in TB in children were mostly female. Based on the Indonesian Health Profile, the number of TB cases in Indonesia in 2019 was 543,874 cases. Compared with

women, the number of TB cases in men is 1.4 times higher in all provinces. Men are more susceptible to TB than women because men have higher mobility and activity than women, so they are more likely to be exposed. This is associated with psychological factors, emotions, and growth and development. There is a tendency to be more active than girls than boys. The risk of exposure to tuberculosis germs is greater in boys who play a lot or do activities outside the home, making it difficult to control. TB cases in men have a risk of 1,311 times compared to women if it is associated with smoking habits in men. In almost all developing countries, smoking prevalence is more than 50% in men compared to women. People who smoke can lower the body's immunity to be more susceptible to TB.

The majority of respondents have normal nutrition, followed by underweight, then obesity, after that several underweight and overweight have the same value. This study is in line with Windy Rakhmawati et al. (2011), which stated that from 74 respondents, 58 children experienced good nutritional status. This is different from the research by Putra & Amelia (2013), which stated that the majority of respondents with TB were children with poor nutritional status. The condition of children with poor nutritional status makes it easier for children to be exposed to infectious diseases, including TB disease. Bodyweight is one of the parameters that give an idea of body mass. Body mass affects the changes, such as when exposed to infection. Nutritional status is closely related to individual problems because it is a predisposing factor that can exacerbate infectious diseases. When malnutrition can affect a person's immune system, a decrease in the body's immune system makes it easier for children to get infectious diseases, including TB disease.

The results showed that most respondents were given exclusive breastfeeding for six months. According to Permatasari and Trijati's (2014) research, TB respondents who were not given exclusive breastfeeding were more than those who gave

exclusive breastfeeding. Breast milk functions to increase the body's immunity, fight germs that enter and can reduce serious illnesses. Infants who are not exclusively breastfed have a 9.19 times risk of TB disease than those who are solely breastfed.¹⁵

This description shows that a higher percentage of children have a history of contact, with 34.3% being household contacts. Children that has history of contact with adults has the main risk factor for children to get TB.¹⁰ Usually, children can become infected with TB after being exposed to adults who have pulmonary TB with a smear (+) result. This study indicates that the percentage of children with a history of TB contacts is smaller when viewed in Nurwitasari's study, which concluded that 87.5% of 24 children.²¹ However, the number of children with a history of contact is quite large, and the results are almost the same as other previous studies, namely the study of Putra and Amelia, which said that there were 30 out of 50 children (60%) who had a history of TB contact.²⁰ In contrast to the Noviarisa study, which showed that only a few children had a history of contact, about 43.9% of the 198 children.²²

In this study, there were more positive scars than negative scars, in contrast to the research of Rosandali et al. (2016). They found that children with TB had more negative scars than positive scars, and there was a significant relationship between the incidence of tuberculosis infection and the BCG vaccine scar formation.⁷ Scar formation after BCG vaccination indicates successful vaccination.²³ After BCG vaccination, children who have scars have better survival overall than children who do not have scars after vaccination.⁴ The effectiveness of BCG vaccination varies between 0-80%.²⁴ The evidence for the protective ability of BCG against childhood TB disease is not very consistent. Still, the results are quite good, ranging from 60-80%, both in developing and developed countries, for pulmonary TB and extrapulmonary TB, although higher levels of protection are found low in the tropics.¹⁰ Several factors affect the scar not forming,

such as incorrect injection method/technique, lack of injection dose, poor vaccine condition/quality, immunity and sensitivity of the child's skin, and the reaction of the child's immune system.²⁵

Parent Characteristic

Most of the patient's parents' last education was higher education, including senior high school and graduate degrees. According to Permatasari and Trijati's (2014) research, more respondents with higher education than those with low education stated no relationship between parental education and children's TB.¹⁵ In contrast to Putri et al. (2015), respondents with low education were more than those with higher education, which confidence a relationship between parental education and children's TB. Low education can affect the level of understanding of child care, such as care, feeding, and guidance to children, which impacts declining health.²⁶ The higher the education of parents, the easier it is to give an understanding of information to maintain the health of children. As in the case of TB, namely by monitoring nutritional intake, BCG vaccination, preventing contact with adult TB sufferers, and paying attention to and maintaining the home environment.⁹

It can be seen from the table of characteristics of parents by occupation that more mothers have the status of IRT. There are similar results from two previous studies, namely the research of Fauza and Riani, which said that more mothers did not have a job or were housewives (63.6%) compared to working mothers.^{27,28}

One of the risk factors that influence the incidence of TB in children is the parents' work status, especially if the child's mother is busy with her work. According to previous research, this is generally because working mothers have little time to spend with their children, so there will be less attention to children's health. In addition, if parents entrust their children to someone else, they will not realize that the person caring for their child has pulmonary TB.¹⁵ In this study, during the interview, several respondents said that the

respondents did not know who their children were in contact with when they were working.

Fathers of TB patients with children have more labor jobs, followed by self-employed and employee jobs. According to Putri et al. (2015), more respondents work than respondents who do not work. Parents' occupations are related to family income. Parents who do not work or whose income is not large can result in the child's nutritional status being less so that they are vulnerable to various diseases, one of which is TB. If the income of a large family is sufficient to meet the nutritional needs of children and children's nutrition can be guaranteed.²⁶

Family income obtained a higher percentage of respondents who earn > UMP. Economic factors are closely related to housing conditions, overcrowding, sanitation, and poor environment increasing TB transmission.²⁹ Lack of family income is also closely related to lack of nutritional intake, which can later lead to weak immunity. In addition, poor economic conditions can lead to a lack of education about TB disease and a lack of access to health services acceptable to patients. A study says that 78.8% (156 respondents) have low socioeconomic status. However, according to research by Permatasari & Trijati (2014), there is no significant relationship between economic status and the incidence of pulmonary TB in children under five.¹⁵

According to research by Samuel (2016), one of the important role in healing and general well-being from TB germ infection is nutrition. However, someone with a low economic status will suffer from high levels of malnutrition due to inadequate nutrition, which in addition to being a risk factor for developing TB disease, can also affect the outcome of TB treatment.³⁰

From this description, parents of children with TB patients with a history of smoking were more than those without a history of smoking. This study is in line with Puspitasari et al. (2015), which stated a history of smoking in parents of TB children in as many as 46 respondents, while there was no history of smoking in parents of TB

children in as many as 40 respondents. Children with smoking parents are said to be passive smokers. Secondhand smoke has a relationship with the development of TB disease. The effect can reduce lung function. Substances contained in cigarettes can impact immunity and body defense mechanisms.³¹

Factors associated with Scar BCG

Based on the results obtained from this study, children with negative scars in children with a history of non-home contact were more than household contacts, and there were no negative scars in children with unknown contacts. Children with positive scars were found in all children with unknown contact history and more than child with household and non-home contacts. There is a significant relationship between the history of contact with BCG scars. This shows the lack of vigilance of parents in controlling or caring for children so that they are exposed to TB sufferers. The parents assume that the scar formed due to the BCG vaccine indicates that the child will be immune to the infection.

In this study, age, gender, nutritional status, history of exclusive breastfeeding, immunization status, mother's occupation, father's occupation, family income, mother's education, and smoking history were not significantly related to BCG scars. A study found that the median scar size in the UK population is bigger than the scar size of BCG vaccination in the Malawian population that appeared three months after.³² In the study of Dhanawade et al., BCG scar formation was not significantly associated with age or gender. The age at which the vaccine was administered is also not always determining factor in scar occurrence.³³ The results of this study are in line with previous studies, which stated that gender and nutritional status of children were not associated with BCG scars.³⁴ In the study of Amriddin et al. (2019), it was shown that there was a significant relationship between socioeconomic status and the nutritional status of infants at birth with BCG scars after controlling for leptin levels.³⁵ BCG scars are a sign of better survival among children in

countries with high child mortality. BCG vaccination could affect the response to several major infections, one of that also includes malaria.³⁶

The limitation of this research is the data collection process using a questionnaire, so there is still the possibility that the respondents did not answer correctly. And the COVID-19 pandemic has prevented researchers from meeting with direct visits to patients' homes, so primary data was taken by telephone.

CONCLUSION

Contact history related to BCG scar in children. It is recommended for parents to pay more attention to and protect their children from the people around them to cut off the possibility of contact with TB patients even though the child has been vaccinated. Medical personnel should supervise/control the quality of the vaccine and the method/technique of injection. And see the scar marks as a successful BCG vaccination.

REFERENCES

1. WHO. Global Tuberculosis Report 2020. 2020.
2. Kemenkes RI. Profil Kesehatan Indonesia Tahun 2019. Jakarta: Kementerian Kesehatan RI; 2020.
3. Ikatan Dokter Anak Indonesia (IDAI). Skar BCG. Jakarta: Ikatan Dokter Anak Indonesia (IDAI); 2016.
4. Benn CS, Roth A, Garly M-L, Fisker AB, Scholtz-Buchholzer F, Timmermann A, et al. BCG scarring and improved child survival: a combined analysis of studies of BCG scarring. *J Intern Med.* 2020;288(6):614–624.
5. Farsida, Hatta M, Patellongi I, Prihantono, Shabariyah R, Larasati Laras RA, et al. The correlation of Foxp3 + gene and regulatory T cells with scar BCG formation among children with Tuberculosis. *J Clin Tuberc other Mycobact Dis.* 2020;21:1–7.
6. Funch KM, Thysen SM, Rodrigues A, Martins CL, Aaby P, Benn CS, et al. Determinants of BCG scarification among children in rural Guinea-Bissau: A prospective cohort study. *Hum Vaccines Immunother.* 2018;14(10):2434–1432.
7. Rosandali F, Aziz R, Suharti N. Hubungan antara Pembentukan Scar Vaksin BCG dan Kejadian Infeksi Tuberculosis. *J Kesehat Andalas.* 2016;5(2):381–384.
8. Sugiyono. Metode Penelitian Kuantitatif dan Kualitatif. Jakarta: Alfabeta; 2012.
9. Brajadenta GS, Laksana ASD, Peramiarti IDSAP. Faktor Risiko Tuberculosis Paru Anak: Studi pada Balai Kesehatan Paru Masyarakat (BKPM) Purwokerto. *Str J Ilm Kesehat.* 2018;7(2):1–6.
10. Rahajoe NN, Supriyatno B, Setyanto DB. Buku Ajar Respirologi Anak. 4th ed. Jakarta: Ikatan Dokter Anak Indonesia; 2018.
11. Farsida, Shabariah R, Hatta M, Patellongi I, Prihantono, Nasrum Massi M, et al. Relationship between expression mRNA gene Treg, Treg, CD4+, and CD8+ protein levels with TST in tuberculosis children: A nested case-control. *Ann Med Surg.* 2020;61:44–47.
12. Kemenkes RI. TB Anak. Jakarta: Kementerian Kesehatan RI; 2019.
13. Rakhmawati FJ, Yulianti AB, Widayanti. Angka Kejadian Tuberculosis Paru pada Anak dengan Imunisasi BCG di RSUD Al-Ihsan Bandung Bulan Januari – Juni 2019. *J Integr Kesehat dan Sains.* 2020;2(2):114–117.
14. Pradani SA, Kundarto W. Evaluasi Ketepatan Obat dan Dosis Obat Anti Tuberculosis pada Pasien Anak Di Instalasi Rawat Jalan RSUD Dr. Moewardi Surakarta Periode 2016-2017. *J Pharm Sci Clin Res.* 2018;3(2):93–103.
15. Permatasari TO, Trijati MH. Karakteristik Individu yang Berhubungan dengan Kejadian Tuberculosis Paru Balita di Balai Kesehatan Paru Masyarakat (BKPM) Kota Cirebon. *Tunas Med J Kedokt Kesehat.* 2014;1(2):1–8.
16. Nelson LJ, Wells CD. Global epidemiology of childhood tuberculosis.

- Int J Tuberc lung Dis. 2004;8(5):636–647.
17. Ridwan H, Rosita Y, Sahfitri A. Faktor Risiko yang Berhubungan dengan Kejadian Tuberkulosis di RSK Paru Palembang Periode Januari-Desember 2010. *Syifa'MEDIKA*. 2012;2(2):98–107.
 18. Dotulong JFJ, Sapulete MR, Kandou GD. Hubungan Faktor Risiko Umur, Jenis Kelamin Dan Kepadatan Hunian Dengan Kejadian Penyakit Tb Paru Di Desa Wori Kecamatan Wori. *J Kedokt Komunitas Dan Trop*. 2015;3(2):57–65.
 19. Rakhmawati W, Fatimah S, Trisyani M, Nurhidayah I. Faktor-Faktor yang Berhubungan dengan Kejadian Tuberkulosis pada Anak di Kecamatan Ngamprah Kabupaten Bandung Barat. *Maj Keperawatan Unpad*. 2011;13(1).
 20. Putra IA, Amelia. Profil Tuberkulosis Pada Anak di Instalasi Rawat Jalan RSUD Raden Mattaher Jambi. *Jambi Med J*. 2013;1:51–60.
 21. Nurwitasari A, Wahyuni CU. Pengaruh Status Gizi dan Riwayat Kontak terhadap Kejadian Tuberkulosis Anak di Kabupaten Jember. *Berk Epidemiol*. 2015;3(2):158–69.
 22. Noviarisa N, Yani FF, Basir D. Tren Kasus Tuberkulosis Anak di RSUP Dr. M. Djamil Padang Tahun 2014-2016. *J Kesehat Andalas*. 2019;8 (Supplement 1):36–41.
 23. Birk NM, Nissen TN, Ladekarl M, Zingmark V, Kjergaard J, Jensen TM, et al. The association between Bacillus Calmette-Guérin vaccination (1331 SSI) skin reaction and subsequent scar development in infants. *BMC Infect Dis*. 2017;17(1):1–8.
 24. Li J, Zhao A, Tang J, Wang G, Shi Y, Zhan L, et al. Tuberculosis vaccine development: from classic to clinical candidates. *Eur J Clin Microbiol Infect Dis*. 2020;39(8):1405–25.
 25. Tabatabaei H. Newborn BCG Vaccinations: Scar Formation and Tuberculin Conversion Rates: A Retrospective Study. *Am J Pediatr*. 2019;5(1):1–6.
 26. Putri RF, Sulastrri D, Lestari Y. Faktor-Faktor yang Berhubungan dengan Status Gizi Anak Balita di Wilayah Kerja Puskesmas Nanggalo Padang. *J Kesehat Andalas*. 2015;4(1):254–261.
 27. Fauza R. Pengaruh Status Gizi Terhadap Kejadian Tb Paru Pada Anak Usia 1-5 Tahun Yang Telah Mendapatkan Imunisasi Bcg Di Rsu Imelda Pekerja Indonesia Tahun 2016. *J Ilm Kebidanan Imelda*. 2019;5(2):68–73.
 28. Riani RES, Machmud PB. Kasus Kontrol Hubungan Imunisasi BCG dengan kejadian TB Paru pada anak tahun 2015-2016. *Sari Pediatr*. 2018;19(6):321–7.
 29. Manalu HSP. Faktor-Faktor Yang Mempengaruhi Kejadian TB Paru Dan Upaya Penanggulangannya. *J Ekol Kesehat*. 2010;9(4):1340–1346.
 30. Samuel B, Volkmann T, Cornelius S, Mukhopadhyay S, MejoJose, Mitra K, et al. Relationship between Nutritional Support and Tuberculosis Treatment Outcomes in West Bengal, India. *J Tuberc Res*. 2016;4(4):213–219.
 31. Puspitasari R, Saraswati L, Hestningsih R. Faktor Yang Berhubungan Dengan Kejadian Tuberkulosis Pada Anak (Studi Di Balai Kesehatan Paru Masyarakat Semarang). *J Kesehat Masy*. 2015;3(1):191–197.
 32. Lalor MK, Ben-Smith A, Gorak-Stolinska P, Weir RE, Floyd S, Blitz R, et al. Population Differences in Immune Responses to Bacille Calmette-Guérin Vaccination in Infancy. *J Infect Dis*. 2009 Mar;199(6):795–800.
 33. Dhanawade SS, Kumbhar SG, Gore AD, Patil VN. Scar formation and tuberculin conversion following BCG vaccination in infants: A prospective cohort study. *J Fam Med Prim care*. 2015;4(3):384–387.
 34. Atimati A, Osarogiagbon O. Prevalence of BCG scar among BCG-vaccinated children in a southern Nigeria tertiary hospital. *Niger J Paediatr*. 2014;41(3):229–233.
 35. Amaruddin AI, Wahyuni S, Hamid F, Chalid MT, Yazdanbakhsh M, Sartono E.

- BCG scar, socioeconomic and nutritional status: a study of newborns in an urban area of Makassar, Indonesia. *Trop Med Int Heal*. 2019;24(6):736–746.
36. Roth A, Gustafson P, Nhaga A, Djana Q, Poulsen A, Garly ML, et al. BCG vaccination scar associated with better childhood survival in Guinea-Bissau. *Int J Epidemiol*. 2005;34(3):540–575.

ORIGINALITY REPORT

19%

SIMILARITY INDEX

17%

INTERNET SOURCES

15%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	event.ners.unair.ac.id Internet Source	4%
2	Submitted to Universitas Indonesia Student Paper	3%
3	"1st Annual Conference of Midwifery", Walter de Gruyter GmbH, 2020 Publication	3%
4	ejournal.upnvj.ac.id Internet Source	2%
5	www.ncbi.nlm.nih.gov Internet Source	1%
6	fatcat.wiki Internet Source	1%
7	Farsida, Rahmini Shabariah, Mochammad Hatta, Ilhamjaya Patellongi et al. "Relationship between expression mRNA gene Treg, Treg, CD4+, and CD8+ protein levels with TST in tuberculosis children: A nested case-control", Annals of Medicine and Surgery, 2021 Publication	1%

8

Sujata Kapil Murarkar, Jayashree Sachin
Gothankar, Prakash Doke, Prasad Pore et al.

"Prevalence and determinants of
undernutrition among under-five children
residing in urban slums and rural area,
Maharashtra, India: a community-based
cross-sectional study", Research Square, 2020

Publication

<1 %

9

pubmed.ncbi.nlm.nih.gov

Internet Source

<1 %

10

"Enhancing Capacity of Healthcare Scholars
and professionals in Responding to the Global
Health Issues", Walter de Gruyter GmbH, 2019

Publication

<1 %

11

Qonita Rahmadiena, Erika Diana Risanti,
Listiana Masyita Dewi, Shinta Riana Setiawati.

"LOW BIRTH WEIGHT AND IMMUNIZATIONS
STATUS: RISK FACTORS OF ACUTE
RESPIRATORY INFECTION IN CHILDREN 2-5
YEARS", Epidemiology and Society Health
Review (ESHR), 2021

Publication

<1 %

12

academic.oup.com

Internet Source

<1 %

13

worldnutrijournal.org

Internet Source

<1 %

14

ejournalmalahayati.ac.id

Internet Source

<1 %

15

journal.universitaspahlawan.ac.id

Internet Source

<1 %

16

www.turkiyeklinikleri.com

Internet Source

<1 %

17

online-journal.unja.ac.id

Internet Source

<1 %

18

ejournal.unisba.ac.id

Internet Source

<1 %

19

Galuh Chandra Irawan, Ani Margawati, Ali Rosidi. "Underweight increases the risk of pulmonary tuberculosis in adult", *Universa Medicina*, 2017

Publication

<1 %

20

Habteyes Tola, K Holakouie-Naieni, Mohammad Ali Mansournia, Mehdi Yaseri et al. "National treatment outcome and predictors of death and treatment failure in multidrug-resistant tuberculosis in Ethiopia: a 10-year retrospective cohort study", *BMJ Open*, 2021

Publication

<1 %

21

bmcpublichealth.biomedcentral.com

Internet Source

<1 %

e-journal.poltekkesjogja.ac.id

22

Internet Source

<1 %

23

e-journal.unair.ac.id

Internet Source

<1 %

24

eprints.umm.ac.id

Internet Source

<1 %

25

www.tandfonline.com

Internet Source

<1 %

26

Nisaulhusna Nisa Ul husna, Suharmadji Suharmadji, Wulan Sari, Agus Alamsyah, Zulmeriza Rasyid. "Faktor Risiko Kejadian Tuberculosis Paru Di Wilayah Kerja Puskesmas Rumbai Kecamatan Rumbai Pesisir Tahun 2019", Jurnal Kesehatan Komunitas, 2020

Publication

<1 %

27

bmcpediatr.biomedcentral.com

Internet Source

<1 %

28

ejournal.unp.ac.id

Internet Source

<1 %

29

eprints.ums.ac.id

Internet Source

<1 %

30

files.givewell.org

Internet Source

<1 %

31

repository.kertacendekia.ac.id

Internet Source

<1 %

32 vdocuments.mx
Internet Source

<1 %

33 hdl.handle.net
Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On