

---

## THE EFFECT OF HIGH-INTENSITY INTERVAL TRAINING ON BLOOD LACTATE LEVELS AND RATE OF PERCEIVED EXERTION IN SEDENTARY HEALTHY ADULTS

Nur Ahlina Damayanti<sup>1\*</sup>, Nury Nusdwinuringtyas<sup>1</sup>, Tresia Fransiska Uliana Tambunan<sup>1</sup>, Aria Kekalih<sup>2</sup>

<sup>1</sup> Department of Physical Health & Rehabilitation, Faculty of Medicine, Universitas Indonesia

<sup>2</sup> Department of Community Medicine, Faculty of Medicine, Universitas Indonesia

\*Correspondence email: [maya\\_15\\_12@yahoo.com](mailto:maya_15_12@yahoo.com)

---

### ABSTRACT

The high-intensity interval training has been seen to be beneficial for various populations in increasing physical capabilities and health quality. This study aimed to determine the effect of high-intensity interval training (HIIT) on blood lactate levels and rate of perceived exertion (RPE) in sedentary healthy adults to discuss the objective outcome of HIIT. A pre-post study was conducted on adults that were stated healthy and sedentary on physical activity. All participants were given HIIT with a total duration of 20 minutes for 12 sessions using a treadmill. Blood lactate level and RPE was recorded on the 1st and 12th session. A significant decrease in blood lactate level and RPE after giving HIIT was found with a reduction of 1,1 mmol/ L and two on the Borg Scale, respectively ( $p < 0.0001$ ). Further research is needed using field-based exercise testing to determine exercise prescription and to obtain the benefit of HIIT in populations with cardiovascular risk factors before utilizing it in patients.

**Keywords:** High-intensity interval training; HIIT; Lactate; Rate of perceived exertion

---

### INTRODUCTION

High-intensity interval training (HIIT) is a short repeatable period of exercise that uses full effort or with an intensity almost reaching max VO<sub>2</sub> ( $\geq 90\%$  of max VO<sub>2</sub>). The exercise could last from a few seconds to minutes. The repeatable exercise is divided into a few sessions with a short rest period in between.<sup>1,2</sup> This type of exercise will stimulate similar physiological changes to continuous mid-intensity exercise with shorter duration and exercise volume.<sup>3</sup> The metabolic adaptive process will be induced faster in HIIT compared to continuing endurance training. HIIT is also proven to be more beneficial in sedentary individuals than continuous training, increasing max VO<sub>2</sub>, and increasing aerobic fitness.<sup>4</sup> The reduction in lactate level and perceived exertion rate in sedentary individuals is seen as an adaptation to HIIT. Other changes in cellular levels are also observed, such as an increase in capillary density and mitochondria

Density in each muscle fiber, will increase the capacity of free fatty acid transport from plasma to cells' cytoplasm to the mitochondria. Other adaptive responses due to HIIT on mitochondria also include higher oxidative metabolism due to increased oxygen uptake, increased lipid metabolism, reduction of hydrogen ion synthesis, and increased lactate deposition in the liver.

The high-intensity interval training has been seen to be beneficial for various populations. The benefit of HIIT is observed in obese individuals with the increase of lipid oxidation with the increasing oxidation capacity of mitochondria with the protein kinase pathway.<sup>1,5,6</sup> Patients with chronic obstructive pulmonary disease (COPD) and coronary artery disease (CAD) also gained substantial benefits with beneficial effects on peripheral muscle without a high increase of burden to the cardiorespiratory system. Patients with CAD are also observed to have higher stroke volume, myocardial contraction,

and an increase in ejection fraction at the end of the exercise.<sup>7,8</sup> Patients with COPD will also experience increasing oxidative muscle capacity and lactate clearance during the resting period, which can decrease ventilation burden and reduce dyspnoea in patients.<sup>9</sup>

Various variables and parameters are used to evaluate the intensity of training. One of which is plasma lactate level. The plasma lactate level is usually used as an exercise intensity marker to evaluate endurance training and/or predict lactate-based energy expenditure from a certain activity. Lactate is commonly produced by red blood cells, neural tissues, intestinal tissues, and dermal tissues. However, the muscle tissue becomes the main producer of lactate during exercise or training.<sup>10</sup> Other parameters used are perceived exertion, which can represent other factors, including psychological status and fitness status.<sup>11</sup>

The effect of HIIT on sedentary individuals is known to have a beneficial effect, as to improve heart rate variability and postural stability in an elderly population.<sup>12,13</sup> However. This study is intended to objectively observe the effect of high-intensity interval training (HIIT) on plasma lactate level and perceived exertion in sedentary healthy adults. Plasma lactate and perceived exertion also might become a new indicators in evaluating the impact of HIIT on each individual.

## **MATERIAL AND METHODS**

This research used an interventional study with a pre-post design with a targeted population of healthy adults in Indonesia. The research is conducted in the General Hospital of Cipto Mangunkusumo, Jakarta, Indonesia. The trial has been approved for ethical review by the health research ethics committee in the Faculty of Medicine Universitas Indonesia with a registration number of 706/UN2.F1/ETIK/PPM.00.02/2019.

### **Sampling procedure and participant's criteria**

The enrolment process of participants used consecutive sampling out of all healthy

adult individuals in the hospital. The inclusion criteria of participants include adults aged 18-40 years old who volunteered to follow the research protocol and exercise regimen by signing the informed consent form, no cardiorespiratory issues, and low activity level based on the International Physical Activity Questionnaire. The low activity level on IPQA is considered insufficiently active (Category 1 of IPQA) if there are less than three days of vigorous activities, less than five days of moderate activities, or less than 600-MET (Metabolic Equivalents)-min/week as mentioned in the guideline.<sup>14</sup> The participants will be excluded if they could not understand the exercise instruction or research protocol, have a history of cardiorespiratory ailments, have balance system disorder, and have injury or abnormalities on lower extremities. Participants will be declared as dropouts should they refuse or not be able to execute HIIT three times in a row or not reach 80% of heart rate on three consecutive exercises.

### **Trial procedure**

At the beginning of the study, all subjects were briefed on the study protocol, then signed informed consent should they consent to the study. Each subject who had signed informed consent performed anamnesis, filled out a PARQ+ questionnaire for pre-exercise screening, IPAQ to determine activity levels, electrocardiography (ECG) examination, and lung function test examination using a peak flow meter and spirometry to determine the cardiorespiratory status of all participants. This data collection is done by researchers and research assistants who have been briefed to fill out questionnaires and conduct examinations.

Each subject who met the inclusion and exclusion criteria was then made to prepare for the training test, which included changing clothes and sports shoes, vital sign checks, oxygen mask installation, Polar® heart rate monitors, and gas analyzers. The training test in this study uses the CPET Cosmed K5® tool, which is calibrated prior to the use of the tool. The patient first undergoes familiarization to get used to walking on the

treadmill for one minute. The researcher describes the training test process and how the participants should signal if they are about to stop. During the training test, the patient is monitored for heart rate, saturation, and blood pressure until there is an indication of discontinuation of the training test and if the subject's efforts reach the maximum. In the training test, the maximum heart rate will be recorded, which then becomes a reference for a high-intensity interval training program. The subjects then sat in a chair for the recovery phase for 15 minutes and underwent remeasurement of the vital signs.

After underwent a training test and getting the maximum heart rate, then the dose of exercise intensity is calculated by using  $Hr_{reserve} (Hr_{max} / peak - Hr_{rest})$  for the intensity of 80% and 40%, which then becomes the reference for the exercise program. 80% intensity will be set for the high-intensity exercise, while the intensity of 40% will be set for low-intensity exercise, which becomes the interval on a high-intensity interval training program. Exercise is given for 20 minutes with a duration of 60 seconds for high intensity and 120 seconds for low intensity. Before, during, and after the exercise, the participants are monitored for their vital signs and perceived exertion rate. In the first exercise, the participants will be measured for their lactate levels right after the finished regiment, using Accutrend® Plus, and asked about the level of effort felt after underwent the exercise. On the second to eleventh exercise, the participants underwent an exercise program of the same intensity. In the twelfth exercise, another examination is carried out as done in the first exercise (lactate measurement and perceived exertion rate examination).

### Statistical analysis

All retrieved data from the trial will be inputted and displayed in tables and diagrams as needed. The statistical analysis is carried out in sequence, starting with univariate analysis (descriptive analysis) on the participants' characteristics and a paired T-test on both pre-exercise and post-exercise groups

to observe the significance of its comparison. The confidence interval is set at 95%. Thus p-value of  $<0.05$  will be considered significant. All the statistical analyses were conducted using IBM SPSS ver 23.0. All retrieved data from the trial will be inputted and displayed in tables and diagrams as needed. The statistical analysis is carried out in sequence, starting with univariate analysis (descriptive analysis) on the participants' characteristics and a paired T-test on both pre-exercise and post-exercise groups to observe the significance of its comparison. The confidence interval is set to 95%. Thus p-value of  $<0.05$  will be considered significant. All the statistical analyses were conducted using IBM SPSS ver 23.0.

### RESULT

A total of 44 participants were included in this study. Due to the consecutive nature of the study, no selection or blinding is conducted. Out of the 44 participants, there were 28 female participants with 16 males in the study. The characteristics of the participant could be seen in Table 1. The mean age of the participants was 30.1 years old, with a mean BMI of 23.5 kg/m<sup>2</sup>. All the participants are of bachelor's degrees implying a similar understanding of the instruction. Based on the International Physical Activity Questionnaire, the median MET value was 330 (120-594) METs. Other physiological indicators of the participants are also recorded with FEV1 (Forced expiratory volume in 1 second), FVC (Forced vital capacity), PFR (Peak flow rate), and PCF (Peak cough flow) values.

**Table 1.** Participants characteristics

	n (%)	Mean (SD)
Age		30,1 (3,6)
Gender		
Male	16 (36,4)	
Female	28 (63,6)	
Body weight (kg)		62,8 (12,6)
Body height (m)		1,6 (0,1)
Education level		
≥ Bachelor	44 (100)	
Maximum heart rate		176 (12)
RPE		16 (1)
IMT		23,5 (3,3)
IPAQ		330 (120 – 594)*
FEV1		2835 (1830 – 4550)*
FVC		3050 (1980 – 5250)*
PFR		478,8 (90,2)
PCF		469,6 (78,3)

\*median (minimum-maximum)

In the first training session, the median value of lactate level in the participants was 6.8 (3-20) mmol/L, as compared to the 12th session, with a median value of lactate level was 5.5 (3-13) mmol/L (Table 2).

**Table 2.** Lactate level and RPE outcomes in 1<sup>st</sup> and 12<sup>th</sup> session

Outcome parameter	1 <sup>st</sup> exercise session	12 <sup>th</sup> exercise session	p-value*
Lactate level	6.8 (3-20) <sup>b</sup>	5.5 (3-13) <sup>b</sup>	<0.0001
Rate of perceived exertion	12.3 (1.4) <sup>a</sup>	10.5 (6-13) <sup>b</sup>	<0.0001


exertion (Borg scale)

<sup>a</sup>Outcome is reported in mean (SD) if considered to have normal distribution

<sup>b</sup>Outcome is reported in median (min-max) format due to abnormal distribution


\*Wilcoxon Signed Ranked Test; p-value of <0.05 is considered significant

The graph in Figure 1 also reports decreasing mean lactate levels from 8.03 to 6.00 in the 1st and 12th exercise sessions. The comparison of lactate measurement in the first and final session is deemed significant (p-value <0.0001).



**Figure 1.** The difference in lactate levels on the 1<sup>st</sup> and 12<sup>th</sup> sessions. There is a decrease in mean blood lactate levels of 8.03 to 6.00 mmol/L.

Similar results can also be inferred from the physical exertion rate used by participants, with 12.3±1.4 on the first session and 10.5 (6-13) on the final session, with a significant difference (p-value <0.0001), as seen in



**Figure 2.** The difference in the rate of perceived exertion (RPE) on the 1<sup>st</sup> and 12<sup>th</sup> sessions, with a decrease of two scores in the Borg Scale.

## DISCUSSION

Despite not being the first study of the HIIT effect on plasma lactate or RPE, the study might be the first to evaluate the output in sedentary individuals. Our research has revealed the effect of HIIT on sedentary individuals with a significant decrease in blood lactate level of 1.1 mmol/L and a decrease of RPE on the Borg scale with a difference of two. The result has concluded the beneficial effect of HIIT on sedentary individuals.

The study's inclusion criteria are set to the age range of 18 to 40 years, based on the previous research conducted by Cardoso et al. (2012) in Brazil with ages between 18-40 years and based on the definition of young adults by Hurlock (2001).<sup>15</sup> With such criteria, 44 participants of 18-40 years old with a mean age of 30 years old were included. The average body mass index of subjects in this study was 23.5 (3.3) kg/m<sup>2</sup>, which was classified as overweight. Physical inactivity or sedentary has a large contribution to the occurrence of excess weight and obesity.<sup>16</sup> The weight gain itself is a significant risk factor for metabolic and heart disease.<sup>17</sup>

The overall level of education of the subject is a bachelor's degree, which eases the

process of filling out English questionnaires and further assistance in providing explanations for the training to be given as well as motivation to follow the exercise program up to twelve times. Out of all 44 participants, there were no drops out in this study.

In this study, the results of the electrocardiogram record of all participants showed no change from all the recorded leads. The examination of lung function described from spirometry results also stated no signs of obstruction or restrictions adjusted to the value of pneumomobile results. The peak cough flow (PCF) value obtained was of  $469.6 \pm 78.3$  L/minute. This result is in line with previous research which stated the normal susceptibility of peak cough flow in sedentary young adult individuals is between 310 - 645 L/minute.<sup>18</sup> All participant is trained to then get intensity to undergo an exercise program, should all of their examination results (ECG, spirometry, and PCF) be considered normal.

The participants of this study were sedentary individuals, where sedentary itself is defined as the unfulfilled threshold of moderate and vigorous-intensity physical activity.<sup>19-23</sup> A study by Tudor-Locke and Meyer (2001) concluded that sedentarism is a low energy use, both for distance and walking time, climbing stairs, and/or low participation in strenuous-intensity activities for either recreation or exercise.<sup>24</sup> Based on the IPAQ results, All the study participants had low levels of physical activity with a middle value of 330 METs per week and a range between 120 - 594 METs. A person falls into the category of low activity if it does not meet the criteria of at least a total of 600 METs in a week, either light, moderate or heavy activities.<sup>16,25</sup> The statement is also supported as a sedentary lifestyle, according to ACSM and the American Heart Association does not fulfill the recommended physical activity of at least 30 minutes at moderate intensity as much as five days a week or even 20 minutes of strenuous activity as much as three days a week.<sup>26</sup> The causes of the high sedentary numbers of the participants can be reviewed

from various aspects. The use of transportation, work, recreational time, and homework are aspects that can play a role in a person's sedentary behavior.<sup>27</sup>

A widely used training test for individuals with low activity levels or sedentary is the modified Bruce protocol. This protocol has also been considered safe for overweight and obese individuals.<sup>28-30</sup> The duration of the training test conducted is around 10 to 12 minutes. The participants stopped the training test if they had met one of the criteria of achieving the maximum training test, namely if there was no increase in heart rate by increasing workload, blood lactate level after the training test was  $>8.0$  mmol / L, the perceived exertion rate has reached  $>17$  on the rate perceived exertion scale (RPE) 6 - 20, or Respiratory Exchange Ratio (RER)  $\geq 1.10$ . In this study, the average perceived exertion rate by the subjects was  $16 \pm 1$ , which according to ACSM, had reached the maximum training test criteria.

The result of this study has shown a significant reduction in serum lactate levels.

The results are in line with a previous study, which showed changes in blood lactate concentration which were observed in 10 healthy men, not exercising regularly, who had been given exercise at a 90% intensity interspersed with a resting phase. The exercise is divided into five meetings with an ergo cycle; each lasts 20 minutes. After given a 20-minute exercise, there was a noticeable change in lactate from the blood lactate sampling in the 20th minute of the exercise from 18.3 mmol to 10.2 mmol. In another study conducted by Macedo, a similar trend was found in blood lactate concentration after high-intensity exercise with a phase interval of 30 seconds by 120 seconds.<sup>31</sup> The significant decrease in post-workout lactate production might occur due to decreased glycolysis or increased lactate clearance from contracting muscles.<sup>30</sup> The process might be explained with Ca (2<sup>+</sup>) -mediated modulation of glycogenolysis, which supplies substrates to glycolytic during the resting phase. Thus, the inactivation of glycolysis is reflected in the cessation of contractile activity.<sup>32</sup> The

study did not differ in the sex or age group for the analysis. Despite being a limitation, several studies have mentioned that there are no significant differences in blood lactate levels between both sexes.<sup>33,34</sup> Significant differences are only found in the age group after 70 years old.<sup>35</sup>

The rate of perceived exertion is an effective measuring tool for measuring weights and fatigue during exercise.<sup>36</sup> The rate of perceived exertion is closely related to the concepts of exercise intensity, exercise motivation, emotional states, and diseases that can expand the scope of factors that affect the level of the effort itself. In healthy individuals, the definition of perceived exertion emphasizes the subjective response to the intensity of exercise and the sensory experience of the perceived stimulus.<sup>37</sup>

Blood lactate levels and rate of perceived exertion decreased simultaneously during 12 sessions of high-intensity interval training, which was in line with Alkahtani's research.<sup>5</sup> In the study, a significant decrease in blood lactate levels occurred in the ninth exercise, and a decrease in the rate of effort also occurred in the ninth exercise. The decreased lactate levels are associated with decreased perceived exertion after high-intensity interval training reflecting the metabolic adaptations in peripheral tissues. According to Bonnila (2020), a positive correlation is seen between the changes of lactate levels and perceived exertion.<sup>36</sup> Such adaptations include decreased glycogenolysis and decreased lactate accumulation in healthy, untrained individuals undergoing high-intensity exercise.<sup>32</sup> The difference between sex on the rate of perceived exertion might not be seen in this study. However, several studies have contradicting results. RPE is found to be lower in the female group compared to males with higher endurance, as shown with %VO<sub>2</sub> max and estimation time limit.<sup>38</sup> Females are also observed to have a lower respiratory exchange ratio in the maximum lactate state as compared to males.<sup>33</sup> However, Rascon et al. stated there are no significant differences between both sexes.<sup>34</sup>

This study is the first to use laboratory training tests to determine the dose of an exercise program. Therefore, the strength of this study is its objectiveness and accuracy, which can be inferred as the gold standard for dosage determination in an exercise program applied to healthy individuals to improve cardiorespiratory fitness.

However, the study is not without limitations. Laboratory training tests can't be applied daily where there is no CPET tool for dosage determination. The diversity of physical activity levels in all subjects was also not controlled prior to the study. Thus, high levels of lactate post-exercise are also possible due to the high intensity of activities shortly before doing the exercises. One of the most notable limitations is also the lack of a control group in this study, thus, the predisposing factor on the decrease of blood lactate and RPE that might be attributable to the result is not considered. However, the nature of this study to reveal the beneficial effect of HIIT on the population is objectively analyzed.

## CONCLUSION

High-intensity interval training is observed to reduce the blood lactate level and rate of perceived exertion after 12 sessions. Further research is needed to elaborate on the findings of this study, including the need for a field test to implement the training regimen without a CPET device properly. Further research is to evaluate the training effects on people with cardiopulmonary comorbidities. Due to our limitation, we also suggest the need for basic lactate measurement before the training regimen to objectively observe each subject's lactate threshold.

## REFERENCES

- Gibala MJ, Mcgee SL. Metabolic Adaptations to Short-term High-Intensity Interval Training : A Little Pain for a Lot of Gain ? 2008;2.
- Gibala M. Physiological adaptations to low-volume high intensity interval training. *Sport Sci Exch.* 2015;28(139):1–6.
- Gibala MJ, Little JP, Macdonald MJ, Hawley JA. Physiological adaptations to low-volume, high-intensity interval training in health and disease. *J Physiol.* 2012;590(5):1077–84.
- Laursen PB, Jenkins DG. The scientific basis for high-intensity interval training: optimising training programs and maximising performance in highly trained endurance athletes. *Sport Med.* 2002;32(1):53–73.
- Alkahtani SA, King NA, Hills AP, Byrne NM. Effect of interval training intensity on fat oxidation, blood lactate and the rate of perceived exertion in obese men. *Springerplus.* 2013;2(1):1–10.
- Talanian JL, Galloway SDR, Heigenhauser GJF, Bonen A, Spriet LL. Two weeks of high-intensity aerobic interval training increases the capacity for fat oxidation during exercise in women. *J Appl Physiol.* 2006;102(4):1439–47.
- Rognmo Ø, Hetland E, Helgerud J, Hoff J, Slørdahl SA. High intensity aerobic interval exercise is superior to moderate intensity exercise for increasing aerobic capacity in patients with coronary artery disease. *Eur J Cardiovasc Prev Rehabil.* 2004;11(3):216–22.
- Warburton DER, McKenzie DC, Haykowsky MJ, Taylor A, Shoemaker P, Ignaszewski AP, et al. Effectiveness of high-intensity interval training for the rehabilitation of patients with coronary artery disease. *Am J Cardiol.* 2005;95(9):1080–4.
- Kortianou EA, Nasis IG, Spetsioti ST, Daskalakis AM, Vogiatzis I. Effectiveness of Interval Exercise Training in Patients with COPD. *Cardiopulm Phys Ther J.* 2010;21(3):12–9.
- Felippe LC, Ferreira GA, De-Oliveira F, Pires FO, Lima-Silva AE. Arterialized and venous blood lactate concentration difference during different exercise intensities. *J Exerc Sci Fit.* 2017;15(1):22–6.
- Coutts AJ, Rampinini E, Marcora SM, Castagna C, Impellizzeri FM. Heart rate

- and blood lactate correlates of perceived exertion during small-sided soccer games. *J Sci Med Sport*. 2009;12(1):79–84.
12. Alansare A, Alford K, Lee S, Church T, Jung HC. The effects of high-intensity interval training vs. Moderate-intensity continuous training on heart rate variability in physically inactive adults. *Int J Environ Res Public Health*. 2018;15(7):1–10.
  13. Elboim-Gabyzon M, Buxbaum R, Klein R. The effects of high-intensity interval training (HIIT) on fall risk factors in healthy older adults: A systematic review. *Int J Environ Res Public Health*. 2021;18(22).
  14. Wolin KY, Heil DP, Askew S, Matthews CE, Bennett GG. Wolin KY, Heil DP, Askew S, Matthews CE, Bennett GG. Validation of the International Physical Activity Questionnaire-Short among Blacks. *J Phys Act Health*. 2008 Sep;5(5):746-60. *J Phys Act Heal*. 2008;5(5):746–60.
  15. Valenti VE, Souza C, Abreu LC De. Peak expiratory flow values are higher in older and taller healthy male children : An observational study. 2010;(June 2014).
  16. Rosenberg DE, Bull FC, Marshall AL, Sallis JF, Bauman AE. Assessment of Sedentary Behavior With the International Physical Activity Questionnaire. 2008;30–45.
  17. Ding D, Sallis JF, Hovell MF, Du J, Zheng M, He H, et al. Physical activity and sedentary behaviours among rural adults in suixi , china : a cross-sectional study. *Int J Behav Nutr Phys Act*. 2011;8(1):37.
  18. Fauzan A. Faktor-Faktor yang mempengaruhi Nilai Arus Puncak Batuk pada dewasa muda sehat Indonesia. Universitas Indonesia; 2016.
  19. Church TS, Martin CK, Thompson AM, Earnest CP, Mikus CR, Blair SN. Changes in Weight , Waist Circumference and Compensatory Responses with Different Doses of Exercise among Sedentary , Overweight Postmenopausal Women. 2009;4(2):1–11.
  20. Herianto, Dewi CR. Analisis dan profil tingkat kebugaran mahasiswa jurusan teknik mesin dan industri universitas gadjah mada yogyakarta. *J Teknosains*. 2012;2(1):1–7.
  21. Melanson EL, Gozansky WS, Barry DW, Maclean PS, Grunwald GK, Hill JO, et al. When energy balance is maintained , exercise does not induce negative fat balance in lean sedentary , obese sedentary , or lean endurance-trained individuals. 2009;80045:1847–56.
  22. Mullen SP, Olson EA, Phillips SM, Szabo AN, Wójcicki TR, Mailey EL. Measuring enjoyment of physical activity in older adults : invariance of the physical activity enjoyment scale ( paces ) across groups and time. 2011;(September).
  23. Sims ST, Larson JC, Hutchinson F, Michael Y, Martin L. Physical Activity and Body Mass: Changes in Younger versus Older Postmenopausal Women. *Med Sci Sport Exerc*. 2011;(June).
  24. Tudor-locke CE, Myers AM. Challenges and Opportunities for Measuring Physical Activity in Sedentary Adults. 2001;31(2):91–100.
  25. Cleland C, Ferguson S, Ellis G, Hunter RF. Validity of the International Physical Activity Questionnaire ( IPAQ ) for assessing moderate-to-vigorous physical activity and sedentary behaviour of older adults in the United Kingdom. 2018;3:1–12.
  26. American College of Sports Medicine. ACSM's Guidelines for Exercise Testing and Prescription. 10th ed. Riebe D, Ehrman JK, Liguori G, Magal M, editors. Philadelphia: Walters Kluwer; 2018.
  27. Owen N, Sugiyama T, Eakin EE, Gardiner PA, Tremblay MS, Sallis JF. Adults' Sedentary Behavior. *AMEPRE*. 2011;41(2):189–96.
  28. Aparecida E, Belén A, Jose P, Galindo M. What is the most effective exercise protocol to improve cardiovascular fitness in overweight and obese subjects ? †. *J Sport Heal Sci*. 2017;6(4):454–61.
  29. Vanhecke TE, Franklin BA, Miller WM, Adam T, Coleman CJ, Mccullough



- PA. Clinical Investigations Cardiorespiratory Fitness and Sedentary Lifestyle in the Morbidly Obese. 2009;124:121–4.
30. Green H, Tupling R, Roy B, Toole DO, Burnett M, Grant S. Adaptations in skeletal muscle exercise metabolism to a sustained session of heavy intermittent exercise. *Am J Physiol Endocrinol Metab*. 2000;118–26.
31. Macedo DV, Lazarim FL, Oliveira F, Tessuti LS, Hohl R. Is lactate production related to muscular fatigue? A pedagogical proposition using empirical facts. 2018;302–7.
32. Crowther GJ, Kemper WF, Carey MF, Conley KE. Control of glycolysis in contracting skeletal muscle. II. Turning it off. *Am J Physiol Endocrinol Metab* [Internet]. 2002 [cited 2022 Dec 10];282(1). Available from: <https://pubmed.ncbi.nlm.nih.gov/11739086/>
33. Hafen PS, Vehrs PR. Sex-Related Differences in the Maximal Lactate Steady State. *Sport* 2018, Vol 6, Page 154 [Internet]. 2018 Nov 27 [cited 2022 Dec 10];6(4):154. Available from: <https://www.mdpi.com/2075-4663/6/4/154/htm>
34. RASCON J, TRUJILLO E, MORALES-ACUÑA F, GUROVICH AN. Differences between Males and Females in Determining Exercise Intensity. *Int J Exerc Sci* [Internet]. 2020 [cited 2022 Dec 10];13(4):1305. Available from: </pmc/articles/PMC7523896/>
35. Korhonen MT, Suominen H, Mero A. Age and sex differences in blood lactate response to sprint running in elite master athletes. *Can J Appl Physiol* [Internet]. 2005 [cited 2022 Dec 10];30(6):647–65. Available from: <https://pubmed.ncbi.nlm.nih.gov/16485517/>
36. Bonilla DA, Petro JL, Vargas-molina S, Martí F, Carbone L. Comparison of blood lactate and perceived exertion responses in two matched time- under-tension protocols. 2020;1–11.
37. Borg G. Borg's Perceived Exertion And Pain Scales. *Human Kinetics*. 1998.
38. Garcin M, Fleury A, Mille-Hamard L, Billat V. Sex-related differences in ratings of perceived exertion and estimated time limit. *Int J Sports Med* [Internet]. 2005 Oct [cited 2022 Dec 10];26(8):675–81. Available from: <https://pubmed.ncbi.nlm.nih.gov/16158374/>